




Amyloid Stain Kit (Congo Red)

Description:

The Amyloid Stain Kit (Congo Red) is intended for use in the histological visualization of amyloid in tissue sections. Examination under a polarizing microscope results in green birefringence of amyloid.

Amyloid: Erythrocytes: Eosinophil Granules: Nuclei: Red to Pink Light Orange Orange to Red Blue

- Uses/Limitations: Not to be taken internally. For In-Vitro Diagnostic use only. Histological applications. Do <u>not</u> use past expiration date. Use caution when handling these reagents. Non-Sterile.
- Control Tissue: Freshly cut sections containing amyloid. Cut sections 6-12 microns to show smaller amyloid deposits.



Availability/Contents:

| Kit Contents | <u>Volume</u> | <u>Storage</u> |
|--------------------|---------------|----------------|
| Congo Red Solution | 500 ml | 18-25℃ |
| Hematoxylin | 500 ml | 18-25℃ |
| Bluing Reagent | 500 ml | 18-25℃ |
| | | |

| Required but not included: | 95% Ethyl Alcohol |
|----------------------------|--------------------|
| | 100% Ethyl Alcohol |

 Precautions:
 Congo Red Solution is flammable.

 Keep away from open flame. Avoid contact with skin and eyes. Harmful if swallowed.

 Follow all Federal, State, and local regulations regarding disposal.

 Use in chemical fume hood whenever possible.

Procedure (Standard):

- 1. Deparaffinize sections if necessary and hydrate to distilled water.
- 2. Stain slide with Hematoxylin for 5 minutes.
- 3. Rinse slide in tap water.
- 4. Incubate slide in Bluing Reagent for 30 seconds.
- 5. Rinse slide in distilled water.
- 6. Dip slide in 95% alcohol for 5 seconds.
- 7. Stain slide with Congo Red Solution for 20 minutes.

Storage: 18° C

Page 1/2 Revision Date: 03/29/2016





- 8. Dip twice (quickly) in 100% alcohol.
- 9. Dip repeatedly (4-5 dips) in clearing agent, and mount in synthetic resin.

SS1003-VO

References:

- 1. Puchtler, H, et al: On the binding of Congo Red amyloid. J. Histochem. Cytochem. Vol. 10: pages 355-363, 1962.
- 2. Eastwood, H. & Cole, K.R., Staining of amyloid in buffered Congo Red in 50% ethanol. Stain Technology. Vol. 46: pages 208-209, 1971.
- 3. Carson, F.L., Histotechnology; A Self-Instructional Text, 2nd Edition. ASCP Press, Chicago, IL. Pages 117-121, 1996.
- 4. Churukian, C., Improved Puchtler's Congo Red method. J. of Histotechnology. Vol. 23: pages 139-141, 2000.

Storage: 18° C



25° C