

## SS1043-VO

# Twort's Counterstain Kit

**Description:** The Twort's Counterstain Kit is designed for histological procedures to provide contrast for easier

visualization of tissue sections.

Nuclei: Red

Cytoplasm: Light Green

**Uses/Limitations:** Not to be taken internally.

For In-Vitro Diagnostic use only.


Histological applications.

Do not use if reagents become cloudy. Do not use past expiration date. Use caution when handling reagents.

Non-Sterile.

**Control Tissue:** Any well-fixed tissue.

Cytology Smear.



### Availability/Contents:

Kit Contents	<u>Volume</u>	<u>Storage</u>
Twort's Neutral Red (Part A)	125 ml	18-25℃
Twort's Fast Green (Part B)	30 ml	18-25℃
Acetic Alcohol Solution (2%)	250 ml	18-25℃

**Precautions:** Avoid contact with skin and eyes.

Flammable.

Harmful if swallowed.

Follow all Federal, State, and local regulations regarding disposal.

#### Prepare the Following Solution Immediately Before Use.

Working Twort's Counterstain Solution: 30 ml Distilled Water

(Add components in the order listed) 9 ml Twort's Neutral Red (Part A)

2 ml Twort's Fast Green (Part B)

Storage: 18° C 25° C

Page 1/2

Revision Date: 09/01/2015



## SS1043-VO

#### Procedure:

- 1. Deparaffinize sections if necessary and hydrate to distilled water.
- 2. Stain slide in Working Twort's Counterstain for 5 minutes. Use once and discard.
- 3. Rinse quickly in distilled water.
- 4. Dip once in absolute alcohol.
- 5. Decolorize quickly and carefully in Acetic Alcohol Solution until no more red color comes away (a few seconds).
- 6. Dip two times in absolute alcohol.
- 7. Clear, and mount in synthetic resin.

Storage: 18° C 25° C

Page 2/2